arXiv:astro-ph/0111051v1 2 Nov 2001

WHERE THE VERY BRIGHT MATTER IS A STANDARD RULER

B.F. ROUKEMA[1,2,3], G.A. MAMON[4], S. BAJTLIK[2]
[1] *DARC/LUTH, Observatoire de Paris-Meudon, 5, place Jules Janssen, F-92.195, Meudon Cedex, France*
[2] *Nicolaus Copernicus Astronomical Center, ul. Bartycka 18, 00-716 Warsaw, Poland*
[3] *University of Warsaw, Krakowskie Przedmieście 26/28, 00-927 Warsaw, Poland*
[4] *Institut d'Astrophysique de Paris, 98bis Bd Arago, F-75.014 Paris, France*

Abstract. Where's the matter? The answer seems to be: "distributed according to a power spectrum with at least one feature (local maximum) near 120–130 h^{-1} Mpc". Analyses of the Iovino, Clowes & Shaver quasar candidate catalogue at $z \sim 2$ and the 2dF Quasar Survey 10K Release (2QZ-10K) support this claim, which has previously been made both for low redshift survey analyses and for high ($z \sim 3$) redshift surveys will be presented. This feature (i) offers a comoving standard ruler which can lift the matter density–cosmological constant ($\Omega_{\rm m}$–Ω_Λ) degeneracy and (ii) might be due either to baryonic acoustic oscillations or to Planck epoch physics which survives through inflation. (i) The 95% confidence constraint from the 2QZ-10K is $\Omega_{\rm m} = 0.25 \pm 0.15, \Omega_\Lambda = 0.60 \pm 0.35$. This constraint is independent of cosmic microwave background constraints and type Ia supernovae constraints. The only assumptions required are (a) that the Universe satisfies a perturbed Friedmann-Lemaître-Robertson-Walker model with a possibly non-zero cosmological constant, (b) that the density perturbations in this model on large scales ($\gg 10 h^{-1}$Mpc) remain small ("linear") and approximately spatially fixed in comoving coordinates, (c) that the statistics (power spectrum or correlation function) of the perturbations are redshift independent, and (d) that quasar redshifts are cosmological.

"Where's the Matter?"

Clues from several surveys, mostly at low redshifts, a few at high redshifts (see the full list of references in [4]) and the increasingly significant results in analyses of quasar surveys [2, 5, 6, 7] imply that the answer is "distributed according to a power spectrum with at least one feature (local maximum) near 120–130 h^{-1} Mpc".

Because the radial distance-redshift relation and the curvature relation between radial and tangential distances both depend on local cosmological parameters $\Omega_{\rm m}$ (the matter density) and Ω_Λ (the cosmological constant), any fine structure in the statistics of density perturbations (traced by any collapsed objects such as galaxies, clusters of galaxies, quasars) will appear to occur at incorrect comoving length scales if incorrect values of $\Omega_{\rm m}$ and Ω_Λ are assumed when transforming angular positions and redshifts into comoving spatial positions.

Fine structure features will only appear at the correct length scales for the correct values of $\Omega_{\rm m}$ and Ω_Λ.

In [5] it was found that the $z \sim 2$ quasar candidate survey of [3] had a feature at the low redshift value of 130 h^{-1} Mpc, provided that $0.1 < \Omega_{\rm m} < 0.45$ (68% confidence), for any value $0 \leq \Omega_\Lambda \leq 1$.

The initial release of the 2QZ-10K [1] yields much stronger results. The 95% confidence constraint from the 2QZ-10K is $\Omega_{\rm m} = 0.25 \pm 0.15, \Omega_\Lambda = 0.60 \pm 0.35$. In this case, an *a priori* value of the length scale for a feature was *not* chosen; internal consistency of features between sub-samples in the different redshift ranges $0.6 < z < 1.1$, $1.1 < z < 1.6$ and $1.6 < z < 2.2$ was the only requirement. This yielded the constraint on $\Omega_{\rm m}$ and Ω_Λ and the length scale $2L = 244 \pm 17\, h^{-1}$ Mpc, consistent with a harmonic of the scale found in [5] and by many earlier authors.

Acknowledgements. This research has been supported by the Polish Council for Scientific Research Grant KBN 2 P03D 017 19 and has benefited from the Programme jumelage 16 astronomie France/Pologne (CNRS/PAN) of the Ministère de la recherche et de la technologie (France).



Figure 1: Confidence intervals for rejecting the hypothesis that the first local maximum at scales greater than $200\,h^{-1}$ Mpc in the comoving spatial correlation function of the 2QZ-10K quasars occurs at the same comoving scale in all three redshift intervals. It is clear that only one region of the $\Omega_{\rm m}$–Ω_Λ plane is favoured.

References

[1] Croom, S. M., Smith, R. J., Boyle, B. J., Shanks, T., Loaring, N. S., Miller, L., Lewis, I. J., 2001, MNRAS 322, L29 (arXiv:astro-ph/0104095)

[2] Deng Z., Xiaoyang X., Fang L.-Zh., 1994, ApJ 431, 506

[3] Iovino A., Clowes R., Shaver P., 1996, A&AS 119, 265

[4] Kirilova, D. P., Chizhov, M. V., 2000, MNRAS 314, 256

[5] Roukema B. F., Mamon G. A., 2000, A&A 358, 395 (arXiv:astro-ph/9911413)

[6] Roukema B. F., Mamon G. A., 2001, A&A 366, 1 (arXiv:astro-ph/0010511)

[7] Roukema B. F., Mamon G. A., Bajtlik S., 2001, A&A accepted, (arXiv:astro-ph/0106135)